

www.agricultureobserver.com

October 2020

Article No. :6

**Concept and Application of Forensic Entomology in Crime Investigation** 

Sujatha, G. S. and Sagar, D.

ICAR-Indian Agricultural Research Institute, New Delhi-110 012 Corresponding author\*: <u>sujigowda2709@gmail.com</u>

# ABSTRACT

Forensic Entomology is defined as the use of insects and their arthropod relatives that inhabit decomposing remains to aid in legal investigations. This branch of science is primarily used to determine the time since death (Post-mortem interval or PMI) or site of human death based on identification of arthropods collected from or near corpse. Since many insects are associated with the human body after death, they are always a potential source of evidence in case of murder, suicide, rape or suspicious death. Hence, it is also called as "Medico-Criminal Entomology". Insects mostly involved in the forensic investigations are true flies or Diptera. The exact and correct identification of the species is very important in forensic entomology, as any error in the analysis may lead to raising charges against an innocent person with the true perpetrator remaining at large.

# **INTRODUCTION**

Forensic entomology is the study of the application of insects and other arthropods in criminal investigation Insects or arthropods are found in a decomposing vertebrate corpse or carrion. These insect colonizers can be used to estimate the time of death i.e., time interval between death and corpse discovery, also called postmortem index (PMI), movement of the corpse, manner and cause of death and association of suspects at the death scene. This article is aimed at providing an overview to forensic odontologists on the possibilities of using forensic data based on insects and their larvae morphology, growth histories, species distribution and toxic contents in their tissue in criminal investigation.

### History

The first recorded incident of insects use in criminal investigation dates back to 13<sup>th</sup> century in China as described in Sung Tzu's book called *The washing away of wrongs*. When a farmer was found murdered in a field with a sharp weapon, all the suspects directed to the sickles placed on the ground. Only one sickle attracted blow flies to the trace amount of blood hidden to the naked eye which resulted in the confession by the murderer. The first application of forensic entomology in a modern court house was in 18<sup>th</sup> century in France where entomological data was admitted as proof for acquitting the current occupants of the residence from where the skeletonized remains of a child were found. In the 18<sup>th</sup> century Yovanovich and Megnin's evaluation of the insect succession on corpses established the science of forensic entomology.

# Arthropods and their association with postmortem changes of the human body

As soon as death occurs, cells start dying and enzymes start digesting the cells inside out by a process called autolysis and the body starts decomposing. Bacteria present in the gastrointestinal tract start destroying the soft tissue producing liquids and gases like hydrogen sulphide, carbon dioxide, methane, ammonia, sulfur dioxide and hydrogen. The volatile molecules

called apeneumones escaping from the decomposing body attract insects. Researchers are able to isolate the volatile chemicals released at different stages of decomposition of the body. The volatile molecules released during each stage can modify the insect behavior. Based on the studies done by Crag *et al.* (1950) it was found that putrative sulfur-based compounds were responsible for initially attracting the flies to the decomposing carcass but egg laying or oviposition of the flies are induced by ammonium-rich compounds present on the carrion.

According to Smith (1986) four categories of insects can be found on decomposing carrion: i) Necrophagous species feeding on the carrion; ii) Predators and parasites feeding on the necrophagous species: this group also contains schizophagous species which feed on the body first and which become predaceous on the later stages; iii) Omnivorous species feeding on the carrion and other arthropods like ants, wasps and some beetles; iv) Other species like springtails and spiders which use the corpse as an extension of their environment. The first two groups are found to be more important for the purpose of forensic entomology. They are mainly from the species of the order Diptera (flies) and Coleoptera (beetles). The succession waves in which the arthropods colonize the carrion depend on the state of decomposition of the carrion.

Insects mostly involved in the forensic investigations are true flies or Diptera. The predominant species in this order are Calliphoridae (blow flies), Sacrophagidae (flesh flies) and Muscidae (house flies). Calliphoridae (blow flies), Sacrophagidae (flesh flies) may arrive within minutes following death. Muscidae (house flies) delay colonization until the body reaches bloat stages of decomposition. According to the studies done by K. Tullies and M. L Goff on exposed carrion in a tropical rainforest, it was found that the decomposition process was divided into five stages on the basis of physical appearance of carcasses, internal temperatures and characteristic insect populations:

- i. **Fresh stage (1-2 Days):** This stage begins at the moment of death and ends when the bloating of the carcass is observed. Even though autolysis occurs at this stage gross morphological changes do not occur at this point. The estimation of the time of death by entomological data after 24 hrs is more accurate than medical examiner's estimation based on the soft tissue examination. Insects were seen attracted within the first 10 min of death to the carcass but no egg laying (oviposition) will be found during this state. Cellular breakdown occurs during this stage without morphologic alterations. Even though morphological changes and odors are not obvious to humans, the chemicals released from the cellular breakdown attract insects even in this early stage.
- ii. **Bloated stage (2-7 Days):** Putrefaction of the body begins at this stage. Gases produced by the metabolic activities of anaerobic bacteria cause an inflation of the abdomen and the carcass forming a balloon-like appearance during the later part. Arthropod activities combined with the putrefaction processes cause internal temperatures of the carcass to rise. The greatest numbers of adult Diptera were attracted to the carcasses during this stage. By the fourth day, first- and early second-instar or larval stages of Diptera will be present. By the beginning of day 2, several predators of Dipteran larvae can be also recovered from the carcasses.

- iii. Decay stage (5-13 Days): Abdominal wall is penetrated, resulting in the deflation of the carcass and ending the bloated stage, the internal temperature rises to 14 degrees above the ambient temperature followed by a drop signifying the end of the decay stage. Decaying odors are high during increased temperature and drop with a fall in temperature. There is a steady decrease in the weight of the carcass by 10<sup>th</sup> day. There is a conversion of carcass biomass to dipteran larval biomass. The larvae subsequently depart from the carcass to pupate.
- iv. **Post-decay stage (10-23 Days):** The post-decay stage begins when most of the Diptera larvae leave the carcass, leaving behind bones, cartilage, hair, small portions of tissue, and a large amount of wet, viscous material known as byproducts of decay (BOD). The BOD is the major site of arthropod activity during this stage.
- v. **Remains stage (18-90+ Days):** This stage is characterized by bones with little cartilage remaining and the BOD has dried up. The transition from post-decay to remains stage is gradual, with declining adult and larval Diptera populations.

### Steps in estimating the postmortem index with insect larvae

The insect larvae present on the dead body can provide evidence for the estimation of PMI up to one month. Correct species identification is the initial step. Different species differ in their growth rates and maturation. For estimating the PMI, age of the larvae has to be determined. By measuring the length or dry weight of the oldest larvae and comparing it with the reference data, age of the larvae can be estimated. The rate of development of the larvae is dependent on the surrounding ambient temperatures. Each stage of development has its temperature requirement hence each species has its own defined number of accumulated degree days or accumulated degree hours to complete its development. Once the thermal history of the larvae is obtained, it can be compared with temperatures at the death scene and PMI can be estimated. The first-generation adult flies can also be used to determine the age. They can be identified by the shriveled wings, and tiny abdomen with dull grey colour. When an insect colonizing the carrion in a particular area is known, an insect colonizing succession model can also be used to estimate the PMI.

### Using insect data for determining the site of crime

There are reported differences in the species of insects involved with the decomposing corpse in different habitats and environments. A careful examination can reveal species variation, as species associated with one type of habitat present on a corpse is found to be different from those when the corpse is transported after death.

# DNA analysis for species identification

Identifying the correct species is the important initial step for estimating the age of the larvae. Morphological comparison is usually used for species identification which requires special expertise and is often time-consuming. In order to overcome this difficulty species identification can be done by polymerized chain reaction amplification suitable regions of the larvae genomes and comparing it with reference data.

# ISSN 2582-6611 Sujatha and Sagar

#### **Agriculture Observer**

## Entomotoxicology

The larvae of the flies which feed on the carrion can accumulate drugs ingested by the deceased person. Bodies which are in advanced stages of decomposition or which are skeletonized are difficult to examine for toxicological substances. In these instances, the larvae fed on this body can be macerated and analyzed with techniques like thin-layer chromatography, gas chromatography or mass spectrometry. Toxins can influence the stages of development of the larvae. Cocaine and heroin in the carcass can accelerate the larval development. Poisons like malathion in the carrion can delay the insect colonization.

#### **CONCLUSION**

Forensic entomology is an emerging field in forensic sciences, where the insects feeding on corpses are studied. The exact and correct identification of the species is very important in forensic entomology, as any error in the analysis may lead to raising charges against an innocent person with the true perpetrator remaining at large. Forensic Entomology as a science is yet to find its place in the legal proceedings of our country. Under the article 138 of the evidence act of the Indian constitution, any scientific evidence is allowed before the court of justice to prove the case, thus general awareness about this branch of science is required as well as trained forensic Entomologist.

#### REFERENCES

- Catts, E.P. and Goff, M.L. 1992. Forensic entomology in criminal investigations. *Annual Review Entomology*, **37**:253-72.
- Fradella, H.F., Owen, S.S. and Burke, T.W. 2007. Building bridges between criminal justice and forensic sciences to create forensic studies programs. *Journal of Criminal Justice Education*, **18**(2):261-282.
- Hall, R.D., Anderson, P.C. and Clark, D.P. 2006. A case of human myiasis caused by *Phormia* regina (Diptera: Calliphoridae) in Missouri, USA. *Journal of Medical Entomology*, 23:578-579.
- Lord, W.D. and Stevenson, J.R. 2013. Directory of forensic entomologists: Defense pest management information analysis center, Washington, DC: Walter Reed Army Medical Center.
- Polson, C.J. 1969. The scientific aspects of forensic medicine. *Journal of Medical Entomology*, 43:675-712.
- Raju, A.A. 2019. Forensic Entomology- Insects Role in Criminal and Civil Laws. *International Journal of Current Microbiology Science*, **8**(1): 2479-2483.
- Rekalde, E.E., Maite, G. and Luisa, M. 2015. Developing and easy and efficient protocol for the study of different blowfly instars through scanning electron microscopy. *Ciencia forense: Revista aragonesa de medicina legal*, 12: 113–136
- Singh, D., Bharti, M. and Singh, T. 1999. Forensic entomology in the Indian perspective. *Journal* of Punjab Academy of Science, 1:217-220.
- Smith, K.G.V. 1986. An introduction to the immature stages of British flies. *Handbook of Identification of British Institute*, **10**(4):127-128.

- Spitaleri, S., Romano, C., Luise, E., Ginestra, E. and Saravo, L. 2006. Genotyping of human DNA recovered from mosquitoes found on a crime scene. *International congress Series*, 1288: 574-576.
- Sukontason, K.L., Vogtsberger, R.C., Boonchu, N., Chaiwong, T., Sripakdee, D., Ngern-Klun, R., Piangjai, S. and Sukontason, K. 2003. Larval morphology of *Chrysomya nigripes* (Diptera: Calliphoridae) a fly species of forensic importance. *Journal of Medical Entomology*, 42:233-240.
- Sumodan, P.K. 2002. Insect detectives. Resonance, 7(8) 51-58.
- Tilstone, J.W., Savage, A.K. and Clark, A.L. 2006. Forensic Science: An Encyclopedia of History, Methods, and Techniques, 357: 1191-1194.
- Wells, J.D. and Stevens, J.R. 2008. Application of DNA-based methods in forensic Entomology. *Annual Review of Entomology*, **53**: 103-120.